

Integration

COS 323

Numerical Integration Problems

- Basic 1D numerical integration

- Given ability to evaluate $f(x)$ for any x , find $\int_a^b f(x) dx$
- Goal: best accuracy with fewest samples

- Classic problem – even analytic functions not necessarily integrable in closed form

$$G(x) = \int_{-\infty}^x e^{-t^2} dt$$

- Other problems (future lectures):

- Multi-dimensional integration
- Ordinary differential equations
- Partial differential equations


Quadrature

- Sample $f(x)$ at a set of points
- Approximate by a function
- Integrate function

- Alternatives:
 - Fit single function vs. multiple (piecewise)
 - Even vs. uneven spacing


Trapezoidal Rule

- Approximate function by trapezoid




Trapezoidal Rule

$$\int_a^b f(x) dx \approx (b-a) \frac{f(a) + f(b)}{2}$$




Extended Trapezoidal Rule

$$\int_a^b f(x) dx \approx (b-a) \frac{f(a) + f(b)}{2}$$



Divide into segments of width h ,
piecewise trapezoidal approximation



$$\int_a^b f(x) dx \approx h \left(\frac{1}{2} f(a) + f(x_1) + \cdots + f(x_{n-1}) + \frac{1}{2} f(b) \right)$$

Trapezoidal Rule Error Analysis

- How accurate is this approximation?

$$\int_a^b f(x) dx \approx \frac{(b-a)}{2} (f(a) + f(b)) + \mathcal{E}$$

- Start with Taylor series for $f(x)$ around midpoint m

$$f(x) \approx f(m) + (x-m) f'(m) + \frac{1}{2} (x-m)^2 f''(m) + \frac{1}{6} (x-m)^3 f'''(m) + \frac{1}{24} (x-m)^4 f^{(4)}(m) + \dots$$

Trapezoidal Rule Error Analysis

- Expand LHS:

$$\int_a^b f(x) dx \approx (b-a) f(m) + 0 + \frac{1}{24} (b-a)^3 f''(m) + 0 + \frac{1}{1920} (b-a)^5 f^{(4)}(m) + \dots$$

- Expand RHS:

$$\frac{(b-a)}{2} (f(a) + f(b)) + \mathcal{E} = \frac{1}{2} (b-a) \left[2f(m) + 0 + \frac{1}{4} (b-a)^2 f''(m) + 0 + \frac{1}{192} (b-a)^4 f^{(4)}(m) + \dots \right] + \mathcal{E}$$

Trapezoidal Rule Error Analysis


- So,

$$\mathcal{E} = -\frac{1}{12}(b-a)^3 f''(m) - \frac{1}{480}(b-a)^5 f^{(4)}(m) + \dots$$

- In general, error for a *single* segment proportional to h^3
- Error for subdividing entire $a \rightarrow b$ interval proportional to h^2
 - “Cubic local accuracy, quadratic global accuracy”
 - Exact for linear functions
 - Note that only even-power terms in error: h^2, h^4 , etc.


Determining Step Size

- Change in integral when reducing step size is a reasonable guess for accuracy
- For trapezoidal rule, easy to go from $h \rightarrow h/2$ without wasting previous samples



Simpson's Rule

- Approximate integral by parabola through three points



$$\int_a^b f(x) dx \approx \frac{b-a}{6} \left(f(a) + 4 f\left(\frac{a+b}{2}\right) + f(b) \right) + O(h^5)$$

- Better accuracy for same # of evaluations
 - Global error $O(h^4)$, exact for cubic (!) functions
- Higher-order polynomials (Newton-Cotes):
 - Global error $O(h^{k+1})$ for k odd, $O(h^{k+2})$ for k even

Richardson Extrapolation

- Better way of getting higher accuracy for a given # of samples
- Suppose we've evaluated integral for step size h and step size $h/2$ using trapezoidal rule:

$$F_h = F + \alpha h^2 + \beta h^4 + \dots$$


$$F_{h/2} = F + \alpha \left(\frac{h}{2}\right)^2 + \beta \left(\frac{h}{2}\right)^4 + \dots$$

- Then

$$\frac{4}{3}F_{h/2} - \frac{1}{3}F_h = F + O(h^4)$$

Richardson Extrapolation

- This treats the approximation as a function of h and “extrapolates” the result to $h=0$
- Can repeat:



Open Methods


- Trapezoidal rule won't work if function undefined at one of the points where evaluating
 - Most often: function infinite at one endpoint

$$\int_0^1 \frac{dx}{x^2}$$

- Open methods only evaluate function on the *open* interval (i.e., not at endpoints)


Midpoint Rule

- Approximate function by rectangle evaluated at midpoint




Extended Midpoint Rule

$$\int_a^b f(x) dx \approx (b-a) f\left(\frac{a+b}{2}\right)$$



Divide into segments of width h :





$$\int_a^b f(x) dx \approx h \left(f\left(a + \frac{h}{2}\right) + f\left(a + \frac{3h}{2}\right) + \cdots + f\left(b - \frac{h}{2}\right) \right)$$

Midpoint Rule Error Analysis

- Following similar analysis to trapezoidal rule, find that local accuracy is cubic, quadratic global accuracy
 - Surprisingly, leading-order constant is $\frac{1}{2}$ as big!
 - Better than trapezoidal rule with fewer samples...
- Formula suitable for adaptive methods and Richardson extrapolation, but can't halve intervals without wasting samples

Extended / Adaptive Midpoint Rule

- Can cut interval into *thirds*:



Limits at Infinity


- Usual trick: change of variables

$$\int_a^b f(x) dx = \int_{1/a}^{1/b} \frac{1}{t^2} f\left(\frac{1}{t}\right) dt$$

- Works with a, b same sign, one of them infinite
 - Otherwise, split into multiple pieces
- Also requires f to decrease faster than $1/x^2$
 - Else need different change of variables, if possible!

Other Quadrature Rules

- Nonuniform sampling: complexity vs. accuracy
- Clenshaw-Curtis: Chebyshev polynomials
 - Change of variables: $x = \cos \theta$
 - Sample at extrema of polynomials
 - FFT-based algorithm to find weights
- Gaussian quadrature
 - Optimize sampling locations to get highest possible accuracy: $O(h^{2n})$ for n sampling points



Discontinuities

- All the above error analyses assumed nice (continuous, differentiable) functions
- In the presence of a discontinuity, all methods revert to accuracy proportional to h
 - In general, if the k -th order derivative is discontinuous, can do no better than $O(h^{k+1})$
- **Locally-adaptive methods:** do not subdivide all intervals equally, focus on those with large error (estimated from change with a single subdivision)