



Global Illumination

COS 426

Overview



- Direct Illumination
 - Emission at light sources
 - Scattering at surfaces
- Global illumination
 - Shadows
 - Inter-object reflections
 - Rendering equation
 - Recursive ray tracing
 - More advanced ray tracing
 - Radiosity


Kajiya 1986



Greg Ward

Direct Illumination (last lecture)

- For each ray traced from camera
 - Sum radiance reflected from each light



$$I = I_E + K_A I_{AL} + \sum_i \left(K_D (N \cdot L_i) + K_S (V \cdot R_i)^n \right) I_i$$

Example



Red's Dream (Pixar Animation Studios)

Overview



- Direct Illumination
 - Emission at light sources
 - Scattering at surfaces
- Global illumination
 - Shadows
 - Inter-object reflections
 - Rendering equation
 - Recursive ray tracing
 - More advanced ray tracing
 - Radiosity



Greg Ward

Overview



- Direct Illumination
 - Emission at light sources
 - Scattering at surfaces
- Global illumination
 - **Shadows**
 - Inter-object reflections
 - Rendering equation
 - Recursive ray tracing
 - More advanced ray tracing
 - Radiosity




Greg Ward

Shadows




- Hard shadows from point light sources




Shadows

- Hard shadows from point light sources




Shadows

- Hard shadows from point light sources



Shadows

- Hard shadows from point light sources
 - Cast ray towards light; $S_L=0$ if blocked, $S_L=1$ otherwise




$$I = I_E + K_A I_{AL} + \sum_{i \in \text{lights}} \left(K_D (N \cdot L_i) + K_S (V \cdot R_i)^n \right) S_i I_i$$

Shadows




- Soft shadows from area light sources
 - Umbra = fully shadowed
 - Penumbra = partially shadowed



Shadows


- Soft shadows from area light sources
 - Average illumination for M sample rays per light



$$I = \dots + \sum_{i \in \text{AreaLights}} \frac{1}{M} \sum_{j \in \text{samples}} \left(K_D (N \cdot L_i) + K_S (V \cdot R_i)^n \right) S_{ij} I_{ij}$$

Shadows

- Soft shadows from circular area light sources
 - Average illumination for M sample rays per light




$$I = \dots + \sum_{i \in \text{AreaLights}} \frac{1}{M} \sum_{j \in \text{samples}} \left(K_D (N \cdot L_i) + K_S (V \cdot R_i)^n \right) S_{ij} I_{ij}$$

Shadow Term

Shadows




- Soft shadows from circular area light sources
 - Average illumination for M sample rays per light
 - Generate M random sample points on area light (e.g., with rejection sampling)
 - Compute illumination for every sample
 - Average



$$I = \dots + \sum_{i \in \text{AreaLights}} \frac{1}{M} \sum_{j \in \text{samples}} \left(K_D (N \cdot L_i) + K_S (V \cdot R_i)^n \right) S_{ij} I_{ij}$$

Direct Illumination

- Illumination from polygonal area light sources
 - Average illumination for M sample rays per light



$$I = \dots + \sum_{i \in \text{AreaLights}} \frac{1}{M} \sum_{j \in \text{samples}} \left(K_D (N \cdot L_i) + K_S (V \cdot R_i)^n \right) S_{ij} I_{ij}$$

Overview




- Direct Illumination
 - Emission at light sources
 - Scattering at surfaces
- Global illumination
 - Shadows
 - **Inter-object reflections**
 - Rendering equation
 - Recursive ray tracing
 - More advanced ray tracing
 - Radiosity




Greg Ward

Inter-Object Reflection




Inter-Object Reflection

- **Radiance** leaving point x on surface is sum of reflected **irradiance** arriving from other surfaces




Solid Angle

- Angle in radians




- Solid angle in **steradians**



Light Emitted from a Surface


- Power per unit area per unit solid angle – *Radiance* (L)
 - Measured in $\text{W}/\text{m}^2/\text{sr}$



$$L = \frac{d\Phi}{dA d\omega}$$

Rendering Equation [Kajiya 86]


- Compute radiance in outgoing direction by integrating reflections over all incoming directions



$$L_o(x', \bar{\omega}') = L_e(x', \bar{\omega}') + \int_{\Omega} f_r(x', \bar{\omega}, \bar{\omega}') (\bar{\omega} \cdot \bar{n}) L_i(x', \bar{\omega}) d\bar{\omega}$$

Rendering Equation [Kajiya 86]

- Compute radiance in outgoing direction by integrating reflections over all incoming directions



$$L_o(x', \bar{\omega}') = L_e(x', \bar{\omega}') + \int_{\Omega} f_r(x', \bar{\omega}, \bar{\omega}') (\bar{\omega} \cdot \bar{n}) L_i(x', \bar{\omega}) d\bar{\omega}$$

Overview




- Direct Illumination
 - Emission at light sources
 - Scattering at surfaces
- Global illumination
 - Shadows
 - Inter-object reflections
 - Rendering equation
 - Recursive ray tracing
 - More advanced ray tracing
 - Radiosity



Greg Ward

Recursive Ray Tracing


- Assume only significant irradiance is in directions of light sources, specular reflection, and refraction



$$L_o(x', \vec{\omega}') = L_e(x', \vec{\omega}') + \int_{\Omega} f_r(x', \vec{\omega}, \vec{\omega}') (\vec{\omega} \cdot \vec{n}) L_i(x', \vec{\omega}) d\vec{\omega}$$

Recursive Ray Tracing


- Compute radiance in outgoing direction by summing reflections from directions of lights specular reflections, and refractions



$$I = I_E + K_A I_{AL} + \sum_L \left(K_D (N \cdot L_i) + K_S (V \cdot R_i)^n \right) S_L I_L + K_S I_R + K_T I_T$$

Recursive Ray Tracing


- Same as ray casting, but trace secondary rays for specular (mirror) reflection and refraction



$$I = I_E + K_A I_{AL} + \sum_L \left(K_D (N \cdot L_i) + K_S (V \cdot R_i)^n \right) S_L I_L + K_S I_R + K_T I_T$$

Specular Reflection


- Trace secondary ray in direction of mirror reflection
 - Evaluate radiance along secondary ray and include it into illumination model



$$I = I_E + K_A I_{AL} + \sum_L \left(K_D (N \cdot L_i) + K_S (V \cdot R_i)^n \right) S_L I_L + K_S \underline{I_R} + K_T I_T$$

Refraction

- Trace secondary ray in direction of refraction
 - Evaluate radiance along secondary ray and include it into illumination model



$$I = I_E + K_A I_{AL} + \sum_L \left(K_D (N \cdot L_i) + K_S (V \cdot R_i)^n \right) S_L I_L + K_S I_R + K_T I_T$$

Recursive Ray Tracing




- ComputeRadiance is called recursively

```
R3Rgb ComputeRadiance(R3Scene *scene, R3Ray *ray, R3Intersection& hit)
{
    R3Ray specular_ray = SpecularRay(ray, hit);
    R3Ray refractive_ray = RefractiveRay(ray, hit);
    R3Rgb radiance = Phong(scene, ray, hit) +
                    Ks * ComputeRadiance(scene, specular_ray) +
                    Kt * ComputeRadiance(scene, refractive_ray);
    return radiance;
}
```

Recursive Ray Tracing


- Which paths?



Recursive Ray Tracing



- Specular reflection and refraction -- $LD(S|R)*E$



Overview




- Direct Illumination
 - Emission at light sources
 - Scattering at surfaces
- Global illumination
 - Shadows
 - Inter-object reflections
 - Rendering equation
 - Recursive ray tracing
 - More advanced ray tracing
 - Radiosity

Kajiya 1986




Greg Ward

Beyond Recursive Ray Tracing




Distributed Ray Tracing

- Estimate integral for each reflection by sampling incoming directions




$$L_o(x', \bar{\omega}') = L_e(x', \bar{\omega}') + \sum_{\text{samples}} f_r(x', \bar{\omega}, \bar{\omega}') (\bar{\omega} \cdot \bar{n}) L_i(x', \bar{\omega}) d\bar{\omega}$$

Ordinary Ray Tracing vs. Distribution Ray Tracing




Ray tracing




Distributed ray tracing

Monte Carlo Path Tracing


- Estimate integral for each pixel by sampling paths from camera



Ray Tracing vs. Path Tracing



Ray tracing



Path tracing



Photon Mapping

- Trace rays *forward* from light sources (recursively)
- Store hits on surfaces: “photon map”
- Final “gather” pass backwards from camera: still compute direct lighting, but look up indirect lighting in photon map



Henrik Wann Jensen

Overview



- Direct Illumination
 - Emission at light sources
 - Scattering at surfaces
- Global illumination
 - Shadows
 - Inter-object reflections
 - Rendering equation
 - Recursive ray tracing
 - More advanced ray tracing
 - Radiosity



Greg Ward


Radiosity



- Indirect diffuse illumination – LD^*E




Rendering Equation (1)



$$L_o(x', \bar{\omega}') = L_e(x', \bar{\omega}') + \int_{\Omega} f_r(x', \bar{\omega}, \bar{\omega}') (\bar{\omega} \cdot \bar{n}) L_i(x', \bar{\omega}) d\bar{\omega}$$

Rendering Equation (2)



$$G(x, x') = \frac{\cos \Theta'_i \cos \Theta_o}{\|x - x'\|^2}$$

$$L(x' \rightarrow x'') = L_e(x' \rightarrow x'') + \int_S f_r(x \rightarrow x' \rightarrow x'') L(x \rightarrow x') V(x, x') G(x, x') dA$$

Radiosity Equation



$$L(x' \rightarrow x'') = L_e(x' \rightarrow x'') + \int_S f_r(x \rightarrow x' \rightarrow x'') L(x \rightarrow x') V(x, x') G(x, x') dA$$

Assume everything
is Lambertian

$$\rho(x') = f_r(x \rightarrow x' \rightarrow x'') \pi$$

$$L(x') = L_e(x') + \frac{\rho(x')}{\pi} \int_S L(x) V(x, x') G(x, x') dA$$

Convert to
Radiosities

$$B = \int_{\Omega} L_o \cos \theta d\omega$$

$$L = \frac{B}{\pi}$$

$$B(x') = B_e(x') + \frac{\rho(x')}{\pi} \int_S B(x) V(x, x') G(x, x') dA$$




Radiosity Approximation

$$B(x') = B_e(x') + \frac{\rho(x')}{\pi} \int_S B(x) V(x, x') G(x, x') dA$$


Discretize the surfaces
into “elements”

$$B_i = E_i + \rho_i \sum_{j=1}^N B_j F_{ij}$$

where
$$F_{ij} = \frac{1}{A_i} \int_{A_i} \int_{A_j} \frac{V_{ij} \cos \Theta'_i \cos \Theta_o}{\pi r^2} dA_j dA_i$$



Radiosity Approximation





System of Equations

$$B_i = E_i + \rho_i \sum_{j=1}^N B_j F_{ij}$$

$$E_i = B_i - \rho_i \sum_{j=1}^N B_j F_{ij}$$

$$B_i - \rho_i \sum_{j=1}^N B_j F_{ij} = E_i$$


$$\begin{bmatrix} 1 - \rho_1 F_{1,1} & \cdot & \cdot & \cdot & -\rho_1 F_{1,n} \\ -\rho_2 F_{2,1} & 1 - \rho_2 F_{2,2} & \cdot & \cdot & -\rho_2 F_{2,n} \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ -\rho_{n-1} F_{n-1,1} & \cdot & \cdot & \cdot & -\rho_{n-1} F_{n-1,n} \\ -\rho_n F_{n,1} & \cdot & \cdot & \cdot & 1 - \rho_n F_{n,n} \end{bmatrix} \begin{bmatrix} B_1 \\ B_2 \\ \cdot \\ \cdot \\ \cdot \\ B_n \end{bmatrix} = \begin{bmatrix} E_1 \\ E_2 \\ \cdot \\ \cdot \\ \cdot \\ E_n \end{bmatrix}$$

$$\left(1 - \rho_i \sum_{j=1}^N F_{ij} \right) B_i - \rho_i \sum_{j=1}^N F_{ij} B_j = E_i$$

$$B_i A_i = E_i A_i + \rho_i \sum_{j=1}^N F_{ji} B_j A_j$$

← This is an energy balance equation

Compare with Direct Illumination



Radiosity



- Application
 - Interior lighting design
 - LD^*E
- Issues
 - Computing form factors
 - Selecting basis functions for radiosity
 - Solving large linear system of equations
 - Meshing surfaces into elements
 - Rendering images

Summary



- Global illumination
 - Rendering equation
- Solution methods
 - Sampling
 - Ray tracing
 - Distributed ray tracing
 - Monte Carlo path tracing
 - Discretization
 - Radiosity

Photorealistic rendering
with global illumination
is an integration problem