

Graph Theory: Matchings and Hall's Theorem

COS 341 Fall 2004

Definition 1 A matching M in a graph $G(V, E)$ is a subset of the edge set E such that no two edges in M are incident on the same vertex, i.e. if $\{w, x\}, \{y, z\} \in M$, then the vertices w, x, y, z are distinct.

The size of a matching M is the number of edges in M . For a graph $G(V, E)$, a matching of maximum size is called a *maximum* matching.

Definition 2 If M is a matching in a graph G , a vertex v is said to be M -saturated if there is an edge in M incident on v . Vertex v is said to be M -unsaturated if there is no edge in M incident on v .

If $G(V_1, V_2, E)$ is a bipartite graph then a matching M of G that saturates all the vertices in V_1 is called a *complete* matching (also called a *perfect* matching).

When does a bipartite graph have a complete matching? Given a graph, if we wanted to prove that the graph has a complete matching, we can simply give the edges in the matching. On the other hand, how do you prove that a graph has no complete matching? In this note, we state and prove Hall's theorem which gives a necessary and sufficient condition for the existence of a complete matching in a bipartite graph.

Before we state Hall's theorem, we will need some definitions and preliminaries.

Definition 3 Given a matching M in graph G , an M -alternating path (cycle) is a path (cycle) in G whose edges are alternately in M and outside of M (i.e. if an edge of the path is in M , the next edge is outside M and vice versa). An M -alternating path whose end vertices are M -unsaturated is called an M -augmenting path.

Lemma 1 If M is a maximum matching in a graph $G(V, E)$, there can be no M -augmenting paths in G .

Proof: Assume, for contradiction, that there exists an M -augmenting path P . Then we can flip the edges of P to obtain a new matching by removing the edges of $P \cap M$ and adding the edges of $P \cap \bar{M}$. More formally, we set $M' = M \cup (P \cap \bar{M}) \setminus (P \cap M)$. It is easy to verify that M' is indeed a valid matching in G . Further, $|M'| = |M| + 1$. This contradicts the fact that M is a maximum matching. ■

Given a bipartite graph $G(V_1, V_2, E)$, and a subset of vertices $S \subseteq V_1$, the neighborhood $N(S)$ is the subset of vertices of V_2 that are adjacent to some vertex in S , i.e.

$$N(S) = \{v \in V_2 : \exists u \in S, (u, v) \in E\}$$

If a bipartite graph has a complete matching saturating V_1 , it is easy to see that $|S| \leq |N(S)|$ for every subset $S \subseteq V_1$. Amazingly, this condition is also sufficient to guarantee the existence of a complete matching.

Theorem 1 (Hall's Theorem) *Let $G(V_1, V_2, E)$ be a bipartite graph with $|V_1| \leq |V_2|$. Then G has a complete matching saturating every vertex of V_1 iff $|S| \leq |N(S)|$ for every subset $S \subseteq V_1$.*

Proof: First we prove that the condition of the theorem is necessary. If G has a complete matching M and S is any subset of V_1 , every vertex in S is matched by M into a different vertex in $N(S)$, so that $|S| \leq |N(S)|$.

Now we prove that the condition is sufficient. Suppose that $|S| \leq |N(S)|$ for every subset $S \subseteq V_1$. Assume for contradiction that G has no complete matching. Let M be a maximum matching, i.e. a matching that saturates the maximum number of vertices in V_1 . Since M is not complete, there exists an M -unsaturated vertex s in V_1 . Let Z be the set of vertices of G reachable from s by M -alternating paths. Since M is a maximum matching, there are no M -augmenting paths among these (by Lemma 1). Let $S = Z \cap V_1$ and $T = Z \cap V_2$. Then, every vertex of T is matched under M to some vertex of $S - \{s\}$ and every vertex of $S - \{s\}$ is matched under M to some vertex of T . Thus $|T| = |S| - 1$. Also, $T = N(S)$. Thus S is a subset of V_1 such that $|N(S)| = |S| - 1$, giving a contradiction. This proves the reverse direction. ■