


# COS 429: COMPUTER VISON MULTI-VIEW GEOMETRY (1 lecture)


- Epipolar Geometry
- The Essential and Fundamental Matrices
- The 8-Point Algorithm
- Trifocal tensor
- Reading: Chapter 10

Many of the slides in this lecture are courtesy to Prof. J. Ponce


## Reconstruction / Triangulation



## (Binocular) Fusion




# Epipolar Geometry




- Epipolar Plane
- Baseline
- Epipoles
- Epipolar Lines

## Epipolar Constraint




- Potential matches for  $p$  have to lie on the corresponding epipolar line  $l'$ .
- Potential matches for  $p'$  have to lie on the corresponding epipolar line  $l$ .

# Epipolar Constraint: Calibrated Case




$$\overrightarrow{Op} \cdot [\overrightarrow{OO'} \times \overrightarrow{O'p}] = 0 \quad \longrightarrow$$

$$\mathbf{p} \cdot [\mathbf{t} \times (\mathcal{R}\mathbf{p}')] = 0 \quad \text{with} \quad \begin{cases} \mathbf{p} = (u, v, 1)^T \\ \mathbf{p}' = (u', v', 1)^T \\ \mathcal{M} = (\text{Id}, \mathbf{0}) \\ \mathcal{M}' = (\mathcal{R}^T, -\mathcal{R}^T \mathbf{t}) \end{cases}$$



Essential Matrix  
(Longuet-Higgins, 1981)



$$\mathbf{p}^T \mathcal{E} \mathbf{p}' = 0 \quad \text{with} \quad \mathcal{E} = [\mathbf{t}_\times] \mathcal{R}$$

## Properties of the Essential Matrix


- $\mathcal{E} p'$  is the epipolar line associated with  $p'$ .
- $\mathcal{E}^T p$  is the epipolar line associated with  $p$ .
- $\mathcal{E} e' = 0$  and  $\mathcal{E}^T e = 0$ .
- $\mathcal{E}$  is singular.
- $\mathcal{E}$  has two equal non-zero singular values  
(Huang and Faugeras, 1989).

# Epipolar Constraint: Small Motions

To First-Order:

$$\mathcal{R}(\mathbf{a}, \theta) = e^{\theta[\mathbf{a}_\times]} \stackrel{\text{def}}{=} \sum_{i=0}^{+\infty} \frac{1}{i!} (\theta[\mathbf{a}_\times])^i \quad \rightarrow \quad \begin{cases} \mathbf{t} = \delta t \mathbf{v} \\ \mathcal{R} = \text{Id} + \delta t [\boldsymbol{\omega}_\times] \\ \mathbf{p}' = \mathbf{p} + \delta t \dot{\mathbf{p}} \end{cases}$$


$$\mathbf{p}^T \mathcal{E} \mathbf{p}' = 0 \quad \text{with} \quad \mathcal{E} = [\mathbf{t}_\times] \mathcal{R} \quad \rightarrow \quad \mathbf{p}^T [\mathbf{v}_\times] (\text{Id} + \delta t [\boldsymbol{\omega}_\times]) (\mathbf{p} + \delta t \dot{\mathbf{p}}) = 0$$



$$\mathbf{p}^T ([\mathbf{v}_\times] [\boldsymbol{\omega}_\times]) \mathbf{p} - (\mathbf{p} \times \dot{\mathbf{p}}) \cdot \mathbf{v} = 0$$

Pure translation:  
Focus of Expansion


## Epipolar Constraint: Uncalibrated Case



$$\hat{\mathbf{p}}^T \mathcal{E} \hat{\mathbf{p}}' = 0$$

$$\mathbf{p} = \mathcal{K} \hat{\mathbf{p}} \quad \xrightarrow{\hspace{1cm}} \quad \mathbf{p}^T \mathcal{F} \mathbf{p}' = 0 \quad \text{with} \quad \mathcal{F} = \mathcal{K}^{-T} \mathcal{E} \mathcal{K}'^{-1}$$

$$\mathbf{p}' = \mathcal{K}' \hat{\mathbf{p}}'$$




Fundamental Matrix  
(Faugeras and Luong, 1992)

## Properties of the Fundamental Matrix


- $\mathcal{F} p'$  is the epipolar line associated with  $p'$ .
- $\mathcal{F}^T p$  is the epipolar line associated with  $p$ .
- $\mathcal{F} e' = 0$  and  $\mathcal{F}^T e = 0$ .
- $\mathcal{F}$  is singular.

## The Eight-Point Algorithm (Longuet-Higgins, 1981)

$$(u, v, 1) \begin{pmatrix} F_{11} & F_{12} & F_{13} \\ F_{21} & F_{22} & F_{23} \\ F_{31} & F_{32} & F_{33} \end{pmatrix} \begin{pmatrix} u' \\ v' \\ 1 \end{pmatrix} = 0$$



$$(uu', uv', u, vu', vv', v, u', v', 1) \begin{pmatrix} F_{11} \\ F_{12} \\ F_{13} \\ F_{21} \\ F_{22} \\ F_{23} \\ F_{31} \\ F_{32} \\ F_{33} \end{pmatrix} = 0$$



$$\begin{pmatrix} u_1u'_1 & u_1v'_1 & u_1 & v_1u'_1 & v_1v'_1 & v_1 & u'_1 & v'_1 \\ u_2u'_2 & u_2v'_2 & u_2 & v_2u'_2 & v_2v'_2 & v_2 & u'_2 & v'_2 \\ u_3u'_3 & u_3v'_3 & u_3 & v_3u'_3 & v_3v'_3 & v_3 & u'_3 & v'_3 \\ u_4u'_4 & u_4v'_4 & u_4 & v_4u'_4 & v_4v'_4 & v_4 & u'_4 & v'_4 \\ u_5u'_5 & u_5v'_5 & u_5 & v_5u'_5 & v_5v'_5 & v_5 & u'_5 & v'_5 \\ u_6u'_6 & u_6v'_6 & u_6 & v_6u'_6 & v_6v'_6 & v_6 & u'_6 & v'_6 \\ u_7u'_7 & u_7v'_7 & u_7 & v_7u'_7 & v_7v'_7 & v_7 & u'_7 & v'_7 \\ u_8u'_8 & u_8v'_8 & u_8 & v_8u'_8 & v_8v'_8 & v_8 & u'_8 & v'_8 \end{pmatrix} \begin{pmatrix} F_{11} \\ F_{12} \\ F_{13} \\ F_{21} \\ F_{22} \\ F_{23} \\ F_{31} \\ F_{32} \end{pmatrix} = - \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}$$

Minimize:

$$\sum_{i=1}^n (\mathbf{p}_i^T \mathcal{F} \mathbf{p}'_i)^2$$

under the constraint  
 $|\mathcal{F}|^2 = 1$ .

## Non-Linear Least-Squares Approach (Luong et al., 1993)



Minimize

$$\sum_{i=1}^n [d^2(\mathbf{p}_i, \mathcal{F}\mathbf{p}'_i) + d^2(\mathbf{p}'_i, \mathcal{F}^T\mathbf{p}_i)]$$

with respect to the coefficients of  $\mathcal{F}$ , using an appropriate rank-2 parameterization.


## The Normalized Eight-Point Algorithm (Hartley, 1995)

- Center the image data at the origin, and scale it so the mean squared distance between the origin and the data points is 2 pixels:  $q_i = T p_i$  ,  $q'_i = T' p'_i$ .
- Use the eight-point algorithm to compute  $\mathcal{F}$  from the points  $q_i$  and  $q'_i$  .
- Enforce the rank-2 constraint.
- Output  $T^T \mathcal{F} T'$ .




Data courtesy of R. Mohr and B. Boufama.

With normalization




Without normalization




Mean errors:  
1.0pixel  
0.9pixel

Mean errors:  
10.0pixel  
9.1pixel

# Trinocular Epipolar Constraints




# Trinocular Epipolar Constraints



$$\begin{cases} \mathbf{p}_1^T \mathcal{E}_{12} \mathbf{p}_2 = 0 \\ \mathbf{p}_2^T \mathcal{E}_{23} \mathbf{p}_3 = 0 \\ \mathbf{p}_3^T \mathcal{E}_{31} \mathbf{p}_1 = 0 \end{cases} \rightarrow \text{These constraints are not independent!}$$


$$\mathbf{e}_{31}^T \mathcal{E}_{12} \mathbf{e}_{32} = \mathbf{e}_{12}^T \mathcal{E}_{23} \mathbf{e}_{13} = \mathbf{e}_{23}^T \mathcal{E}_{31} \mathbf{e}_{21} = 0$$

# Trinocular Epipolar Constraints: Transfer



$$\begin{cases} \mathbf{p}_1^T \mathcal{E}_{12} \mathbf{p}_2 = 0 \\ \mathbf{p}_2^T \mathcal{E}_{23} \mathbf{p}_3 = 0 \\ \mathbf{p}_3^T \mathcal{E}_{31} \mathbf{p}_1 = 0 \end{cases} \rightarrow \text{Given } \mathbf{p}_1 \text{ and } \mathbf{p}_2, \mathbf{p}_3 \text{ can be computed as the solution of linear equations.}$$

# Trifocal Constraints



$$z\mathbf{p} = \mathcal{M}\mathbf{P} \iff \mathbf{l}^T \mathcal{M}\mathbf{P} = 0 \iff \mathbf{L} \cdot \mathbf{P} = 0 \text{ with } \mathbf{L} = \mathcal{M}^T \mathbf{l}$$

$$\xrightarrow{\quad} \begin{pmatrix} \mathbf{L}_1^T \\ \mathbf{L}_2^T \\ \mathbf{L}_3^T \end{pmatrix} \mathbf{P} = \mathbf{0} \quad \xrightarrow{\quad} \text{Rank}(\mathcal{L}) = 2 \quad \text{where} \quad \mathcal{L} \stackrel{\text{def}}{=} \begin{pmatrix} \mathbf{l}_1^T \mathcal{M}_1 \\ \mathbf{l}_2^T \mathcal{M}_2 \\ \mathbf{l}_3^T \mathcal{M}_3 \end{pmatrix}$$


# Trifocal Constraints

## Calibrated Case


$$\text{Rank}(\mathcal{L}) = 2 \quad \text{where} \quad \mathcal{L} \stackrel{\text{def}}{=} \begin{pmatrix} \mathbf{l}_1^T \mathcal{M}_1 \\ \mathbf{l}_2^T \mathcal{M}_2 \\ \mathbf{l}_3^T \mathcal{M}_3 \end{pmatrix}$$

$\mathcal{G}_1^i = \mathbf{t}_2 \mathbf{R}_3^{iT} - \mathbf{R}_2^i \mathbf{t}_3^T \quad \text{for } i = 1, 2, 3$

and  $\mathcal{R}_j = (\mathbf{R}_j^1 | \mathbf{R}_j^2 | \mathbf{R}_j^3)$  for  $j = 2, 3$



All 3x3 minors  
must be zero!



Pick  $\mathcal{M}_1 = (\text{Id} \quad \mathbf{0})$ ,  $\mathcal{M}_2 = (\mathcal{R}_2 \quad \mathbf{t}_2)$  and  $\mathcal{M}_3 = (\mathcal{R}_3 \quad \mathbf{t}_3)$ .

$$\mathcal{L} = \begin{pmatrix} \mathbf{l}_1^T & 0 \\ \mathbf{l}_2^T \mathcal{R}_2 & \mathbf{l}_2^T \mathbf{t}_2 \\ \mathbf{l}_3^T \mathcal{R}_3 & \mathbf{l}_3^T \mathbf{t}_3 \end{pmatrix} \longrightarrow \mathbf{p}_1^T \begin{pmatrix} \mathbf{l}_2^T \mathcal{G}_1^1 \mathbf{l}_3 \\ \mathbf{l}_2^T \mathcal{G}_1^2 \mathbf{l}_3 \\ \mathbf{l}_2^T \mathcal{G}_1^3 \mathbf{l}_3 \end{pmatrix} = 0$$


$\mathcal{G}_1^1$   
 $\mathcal{G}_1^2$   
 $\mathcal{G}_1^3$

Trifocal Tensor


# Trifocal Constraints

## Uncalibrated Case

$$\text{Rank}(\mathcal{L}) = 2 \quad \text{where} \quad \mathcal{L} \stackrel{\text{def}}{=} \begin{pmatrix} \mathbf{l}_1^T \mathcal{M}_1 \\ \mathbf{l}_2^T \mathcal{M}_2 \\ \mathbf{l}_3^T \mathcal{M}_3 \end{pmatrix}$$




$$\mathcal{L} = \begin{pmatrix} \mathbf{l}_1^T \mathcal{K}_1 & 0 \\ \mathbf{l}_2^T \mathcal{K}_2 \mathcal{R}_2 & \mathbf{l}_2^T \mathcal{K}_2 \mathbf{t}_2 \\ \mathbf{l}_3^T \mathcal{K}_3 \mathcal{R}_3 & \mathbf{l}_3^T \mathcal{K}_3 \mathbf{t}_3 \end{pmatrix}$$



Pick  $\mathcal{M}_1 = (\mathcal{K}_1 \quad \mathbf{0})$ ,  $\mathcal{M}_2 = (\mathcal{A}_2 \mathcal{K}_1 \quad \mathbf{a}_2)$  and  $\mathcal{M}_3 = (\mathcal{A}_3 \mathcal{K}_1 \quad \mathbf{a}_3)$ .

$$\mathbf{l}_1 \propto \begin{pmatrix} \mathbf{l}_2^T \mathcal{G}_1^1 \mathbf{l}_3 \\ \mathbf{l}_2^T \mathcal{G}_1^2 \mathbf{l}_3 \\ \mathbf{l}_2^T \mathcal{G}_1^3 \mathbf{l}_3 \end{pmatrix} \quad \Rightarrow \quad \mathbf{p}_1^T \begin{pmatrix} \mathbf{l}_2^T \mathcal{G}_1^1 \mathbf{l}_3 \\ \mathbf{l}_2^T \mathcal{G}_1^2 \mathbf{l}_3 \\ \mathbf{l}_2^T \mathcal{G}_1^3 \mathbf{l}_3 \end{pmatrix} = 0$$



$$\begin{matrix} \mathcal{G}_1^1 \\ \mathcal{G}_1^2 \\ \mathcal{G}_1^3 \end{matrix}$$

Trifocal Tensor


## Properties of the Trifocal Tensor

- For any matching epipolar lines,  $\mathbf{l}_2^T \mathcal{G}_1^i \mathbf{l}_3 = 0$ .
- The matrices  $\mathcal{G}_1^i$  are singular.
- They satisfy 8 independent constraints in the uncalibrated case (Faugeras and Mourrain, 1995).

## Estimating the Trifocal Tensor


- Ignore the non-linear constraints and use linear least-squares a posteriori.
- Impose the constraints a posteriori.

For any matching epipolar lines,  $\mathbf{l}_2^T \mathcal{G}_1^i \mathbf{l}_3 = 0$ .



The backprojections of the two lines do not define a line!

## Multiple Views (Faugeras and Mourrain, 1995)



$$z\mathbf{p} = \mathcal{M}\mathbf{P} \iff \mathbf{p} \times (\mathcal{M}\mathbf{P}) = ([\mathbf{p}_x]\mathcal{M})\mathbf{P} = 0$$

$$\begin{pmatrix} u\mathcal{M}^3 - \mathcal{M}^1 \\ v\mathcal{M}^3 - \mathcal{M}^2 \end{pmatrix} \mathbf{P} = 0 \quad \text{where} \quad \mathcal{M} = \begin{pmatrix} \mathcal{M}^1 \\ \mathcal{M}^2 \\ \mathcal{M}^3 \end{pmatrix}$$

$$\mathcal{Q}\mathbf{P} = 0 \quad \text{where} \quad \mathcal{Q} \stackrel{\text{def}}{=} \begin{pmatrix} u_1\mathcal{M}_1^3 - \mathcal{M}_1^1 \\ v_1\mathcal{M}_1^3 - \mathcal{M}_1^2 \\ u_2\mathcal{M}_2^3 - \mathcal{M}_2^1 \\ v_2\mathcal{M}_2^3 - \mathcal{M}_2^2 \\ u_3\mathcal{M}_3^3 - \mathcal{M}_3^1 \\ v_3\mathcal{M}_3^3 - \mathcal{M}_3^2 \\ u_4\mathcal{M}_4^3 - \mathcal{M}_4^1 \\ v_4\mathcal{M}_4^3 - \mathcal{M}_4^2 \end{pmatrix} \implies \boxed{\text{Rank}(\mathcal{Q}) \leq 3}$$

## Two Views

$$QP = 0 \quad \text{where} \quad Q \stackrel{\text{def}}{=} \begin{pmatrix} u_1\mathcal{M}_1^3 - \mathcal{M}_1^1 \\ v_1\mathcal{M}_1^3 - \mathcal{M}_1^2 \\ u_2\mathcal{M}_2^3 - \mathcal{M}_2^1 \\ v_2\mathcal{M}_2^3 - \mathcal{M}_2^2 \\ u_3\mathcal{M}_3^3 - \mathcal{M}_3^1 \\ v_3\mathcal{M}_3^3 - \mathcal{M}_3^2 \\ u_4\mathcal{M}_4^3 - \mathcal{M}_4^1 \\ v_4\mathcal{M}_4^3 - \mathcal{M}_4^2 \end{pmatrix} \implies \text{Rank}(Q) \leq 3$$

$$\text{Det} \begin{pmatrix} u_1\mathcal{M}_1^3 - \mathcal{M}_1^1 \\ v_1\mathcal{M}_1^3 - \mathcal{M}_1^2 \\ u_2\mathcal{M}_2^3 - \mathcal{M}_2^1 \\ v_2\mathcal{M}_2^3 - \mathcal{M}_2^2 \end{pmatrix} = 0 \quad \rightarrow \quad \text{Epipolar Constraint}$$

## Three Views

$$QP = 0 \quad \text{where} \quad Q \stackrel{\text{def}}{=} \begin{pmatrix} u_1\mathcal{M}_1^3 - \mathcal{M}_1^1 \\ v_1\mathcal{M}_1^3 - \mathcal{M}_1^2 \\ u_2\mathcal{M}_2^3 - \mathcal{M}_2^1 \\ v_2\mathcal{M}_2^3 - \mathcal{M}_2^2 \\ u_3\mathcal{M}_3^3 - \mathcal{M}_3^1 \\ v_3\mathcal{M}_3^3 - \mathcal{M}_3^2 \\ u_4\mathcal{M}_4^3 - \mathcal{M}_4^1 \\ v_4\mathcal{M}_4^3 - \mathcal{M}_4^2 \end{pmatrix} \implies \text{Rank}(Q) \leq 3$$


$$\text{Det} \begin{pmatrix} u_1\mathcal{M}_1^3 - \mathcal{M}_1^1 \\ v_1\mathcal{M}_1^3 - \mathcal{M}_1^2 \\ u_2\mathcal{M}_2^3 - \mathcal{M}_2^1 \\ v_3\mathcal{M}_3^3 - \mathcal{M}_3^2 \end{pmatrix} = 0 \quad \rightarrow \quad \text{Trifocal Constraint}$$

## Four Views


$$\mathcal{Q}\mathbf{P} = 0 \quad \text{where} \quad \mathcal{Q} \stackrel{\text{def}}{=} \begin{pmatrix} u_1\mathcal{M}_1^3 - \mathcal{M}_1^1 \\ v_1\mathcal{M}_1^3 - \mathcal{M}_1^2 \\ u_2\mathcal{M}_2^3 - \mathcal{M}_2^1 \\ v_2\mathcal{M}_2^3 - \mathcal{M}_2^2 \\ u_3\mathcal{M}_3^3 - \mathcal{M}_3^1 \\ v_3\mathcal{M}_3^3 - \mathcal{M}_3^2 \\ u_4\mathcal{M}_4^3 - \mathcal{M}_4^1 \\ v_4\mathcal{M}_4^3 - \mathcal{M}_4^2 \end{pmatrix} \implies \text{Rank}(\mathcal{Q}) \leq 3$$

$$\text{Det} \begin{pmatrix} v_1\mathcal{M}_1^3 - \mathcal{M}_1^2 \\ u_2\mathcal{M}_2^3 - \mathcal{M}_2^1 \\ v_3\mathcal{M}_3^3 - \mathcal{M}_3^2 \\ v_4\mathcal{M}_4^3 - \mathcal{M}_4^2 \end{pmatrix} = 0 \quad \rightarrow \quad \text{Quadrifocal Constraint} \\ (\text{Triggs, 1995})$$

Geometrically, the four rays must intersect in  $P..$



# Quadrifocal Tensor and Lines



$$z\mathbf{p} = \mathcal{M}\mathbf{P} \iff \mathbf{l}^T \mathcal{M}\mathbf{P} = 0 \iff \mathbf{L} \cdot \mathbf{P} = 0 \text{ with } \mathbf{L} = \mathcal{M}^T \mathbf{l}$$

$\begin{pmatrix} \mathbf{L}_1^T \\ \mathbf{L}_2^T \\ \mathbf{L}_3^T \\ \mathbf{L}_4^T \end{pmatrix} \mathbf{P} = \mathbf{0}$  Rank( $\mathcal{L}$ ) = 3 where  $\mathcal{L} \stackrel{\text{def}}{=} \begin{pmatrix} \mathbf{l}_1^T \mathcal{M}_1 \\ \mathbf{l}_2^T \mathcal{M}_2 \\ \mathbf{l}_3^T \mathcal{M}_3 \\ \mathbf{l}_4^T \mathcal{M}_4 \end{pmatrix}$

# Scale-Restraint Condition from Photogrammetry

