**COS 511: Foundations of Machine Learning**

Rob Schapire
Scribe: Eric Glover

Lecture #14
March 27, 2003

# 1 Winnow

**review from last time**

$\eta > 0$ ← learning rate
$\vec{w}_{1,i} = 1/N$ ← Initial distribution
for $t = 1, 2, ...T$ ← $T$ steps
  get $\vec{x}_t \in \mathcal{R}^n$
  predict $\hat{y}_t = sign(\vec{w}_t \cdot \vec{x}_t)$ Make prediction for the current step
  observe $y_t \in \{-1, 1\}$
(update:)
if $y_t = \hat{y}_t$ then $\vec{w}_{t+1} = \vec{w}_t$ We got it right, so we don't do any updating
else

$$w_{t+1,i} = w_{t,i} \frac{e^{\eta y_t x_{t,i}}}{Z_t} \tag{1}$$

Equation 1 has the property that if the sign of $y_t x_{t,i}$ is positive, then it will increase $w_{t+1,i}$, and if the sign is negative, it will decrease it.

## 1.1 Analysis

Assume $||\vec{x}_t||_\infty \leq 1$ note: $L_\infty$ norm is the maximum absolute value of any component
$\exists \delta > 0, \vec{u} \in \mathcal{R}^n$ st ← $\vec{u}$ is the true weights
$\forall_t \; y_t(\vec{u} \cdot \vec{x}_t) \geq \delta$ ← for all examples, margin is at least $\delta$
$||\vec{u}||_1 = 1$ ← Sum of the absolute value of all components of $u$ is 1.
$u_i \geq 0$
Thm:

$$\#\text{ mistakes} \leq \frac{\ln N}{\eta\delta + \ln(\frac{2}{e^\eta + e^{-\eta}})} \tag{2}$$

Solving for minimum value for Equation 2, we get:

$$\#\text{ mistakes} \leq \frac{2 \ln N}{\delta^2} \quad \text{if } \eta = \frac{1}{2} \ln(\frac{1+\delta}{1-\delta}) \tag{3}$$

## 1.2 Proof

Measure of progress - how close $\vec{w}_t$ (predicted weights) is to $\vec{u}$ (actual weights).

$\Phi$ = Potential function of measure of progress

Since both $\vec{u}$ and $\vec{w}_t$ are probability distributions, we use Relative Entropy (RE):

$$\Phi_t = RE(\vec{u}||\vec{w}_t) \quad : \quad RE(\vec{p}||\vec{q}) = \sum_i p_i \ln \frac{p_i}{q_i} \tag{4}$$

try to prove every time makes a mistake $\Phi$ drops by some amount. Since RE always $\geq 0$, this gives a bound on the total number of mistakes.

Since nothing happens when the algorithm does not make a mistake, we assume that it makes a mistake on every round.

$$\Phi_{t+1} - \Phi_t = \sum_i u_i ln \frac{u_i}{w_{t+1,i}} - \sum_i u_i ln \frac{u_i}{w_{t,i}} \tag{5}$$

$$\ln(\frac{u_i}{w_{t+1,i}}) = \ln u_i - \ln w_{t+1,i} \qquad and \qquad \ln(\frac{u_i}{w_{t+1,i}}) = \ln u_i - \ln w_{t,i} \tag{6}$$

Given Equation 5 and 6, you get 7:

$$\Phi_{t+1} - \Phi_t = \sum_i u_i \ln \frac{w_{t,i}}{w_{t+1,i}} = \sum_i u_i \ln \frac{Z_t}{e^{\eta y_t x_{t,i}}} \tag{7}$$

$$= \sum_i u_i \ln Z_t - \sum_i u_i \eta y_t x_{t,i} \tag{8}$$

$$= \ln Z_t - \eta y_t (\vec{u} \cdot \vec{x}_t) \tag{9}$$

We know that $y_t(\vec{u} \cdot \vec{x}_t) \geq \delta$ and that

$$Z_t = \sum_i w_i e^{\eta y_t x_{t,i}} \tag{10}$$

So how do we upper bound an exponential?

We upperbound the exponential by a linear as shown in Figure 1.

The new equation using the linear bound is:

$$Z_t \leq \sum_i w_i \left[ \left(1 + \frac{yx_i}{2}\right) e^{\eta} + \left(1 - \frac{yx_i}{2}\right) e^{-\eta} \right] \tag{11}$$

$$\leq \left(\frac{e^{\eta} + e^{-\eta}}{2}\right) \sum_i w_i + \left(\frac{e^{\eta} + e^{-\eta}}{2}\right) \sum_i w_i y x_i \tag{12}$$




Figure 1: Upperbound an exponential on the range [-1,1] by a linear.

Since: $\sum_i w_i = 1$, $\frac{e^{\eta}+e^{-\eta}}{2} > 0$ and since we made a mistake, $y_t(\vec{w}_t \cdot \vec{x}_t) \leq 0$, we can conclude that right half is always negative, and hence the bound from Equation 12 is:

$$Z_t \leq \frac{e^{\eta} + e^{-\eta}}{2} \tag{13}$$

Thus:

$$\Phi_{t+1} - \Phi_t \leq \ln\left(\frac{e^{\eta} + e^{-\eta}}{2}\right) - \eta\delta \tag{14}$$

We define: $c = \ln\left(\frac{e^{\eta}+e^{-\eta}}{2}\right) - \eta\delta$

Continuing:

$$\Phi_1 = \mathrm{RE}(\vec{u}||\vec{w}) = \sum_i u_i \ln(u_i N) \leq \sum_i u_i \ln N = \ln N \tag{15}$$

Thus the first round: $\Phi_1$ has an upperbound of $\ln N$, and each additional round this value must drop by $c$, as shown by Equation 14

Hence, the maximum number of mistakes is: $\leq \frac{\ln N}{c}$.

If $\eta = \frac{1}{2}\ln(\frac{1+\delta}{1-\delta})$ then: $c = \mathrm{RE}(\frac{1}{2}$ - $\frac{\delta}{2}||\frac{1}{2})$ which $\geq 2(\frac{\delta}{2})^2 = \frac{\delta^2}{2}$

Summary:

For perceptron: $\frac{1}{\delta^2} \rightarrow Nk$ mistakes for $k$ experts.

For Winnow: $\rightarrow 2k^2 \ln N$, which is better when $k << N$.

### 1.3 What about the constraint $u_i \geq 0$

Until now, we assumed that $\vec{u}$ is all positive, so how do we permit components of $\vec{u}$ to be negative, or to correspond with negative values, without causing math problems later?

The solution is to duplicate the components of $\vec{x}$, but make the right half (the duplicates) negative, and to have $\vec{u}$ broken into two halves, one for the positive components, and one for the negative components.

For example: lets say we wanted the following:

$\vec{x} = (1, .7, -.4)$ $\vec{u} = (.5, .2, -.3)$

We would duplicate and invert the sign of the components of $\vec{x}$, so:

$\vec{x} = (1, .7, -.4) \rightarrow (1, .7, .-4, \quad -1, -.7, .4)$

For $\vec{u}$ we zero out the negative components on the left, and zero out the positive components on the right as shown:

$\vec{u} = (.5, .2, -.3) \rightarrow (.5, .2, 0 \quad 0, 0, .3)$

This results in the same dot product as if you used your original values for $\vec{u}$ and $\vec{x}$. The resulting algorithm is called the "balanced winnow" algorithm, and is accomplished by doubling the number of weights as described above.

## 2 Estimating Probabilities of Predictions

Previous classification learning problems the goal was to minimize the probability of making a mistake. The question is how do we estimate the probability of a given prediction.

For example:

$x$ is the current weather conditions, and $y$ is the prediction for tomorrow.

$$y = \begin{cases} 1 & \text{if rain tomorrow} \\ 0 & \text{otherwise} \end{cases}$$

This problem is a distribution of pairs $(x, y) \sim D$. The goal is to learn to estimate a distribution: $p(x) = \Pr[y = 1|x]$. This is equal to the expectation or $E[y|x]$. In this case $y$ is binary, although in other problems, $y$ might be a real. For example, $y$ could be the amount of rain on a given day.

We define $h(x)$ as an estimate of $p(x)$ from a given expert. We want $h(x) \approx p(x)$, but we never see $p(x)$, we only see the $x$ values. In otherwords, there might be a 80% chance of rain, although it might not actually rain. All we know is that it didn't rain, not that there was an 80% chance of it.

The method is to penalize $h$ on $(x, y)$ as follows:

$(h(x) - y)^2$ is a loss function, also called a cost function, in this case, square loss, quadratic loss or Breir score.

We have a set of predictions and$(x_1, y_1), \ldots, (x_m, y_m)$ and the actual events. We wish to choose $h$ that minimizes the loss function, as in Equation 16:

$$\sum_i (h(x_i) - y_i)^2 \tag{16}$$

If $h$ is unrestricted, when is the expected loss $E[(h(x)-y)^2]$ minimized? Fix $x$. Let $p = p(x) = \Pr[y = 1], h = h(x)$. Then

$$E[(h - y)^2] = p(h - 1)^2 + (1 - p)h^2 \tag{17}$$

We now minimize over $h$ by taking the derivative with respect to h, and set equal to 0:

$$\frac{d}{dh} = 2p(h - 1) + 2(1 - p)h = 2(h - p) \tag{18}$$

Equation 18 has a minimum when $h = p$. Hence, **the loss function is minimized when h=p**.

Continuing:

$$E_x[\underbrace{(h(x)-p(x))^2}_{\text{goal}}] = E_{x,y}[\underbrace{(h(x)-y)^2}_{\text{observed}}] - E_{x,y}[\quad \underbrace{(p(x)-y)^2}_{\text{Intrinsic randomness}} \quad] \tag{19}$$

Note: the expectaion is over both $x, y$, since it is constant in terms of h. Also, the $p(x)$ is the intrinsic randomness, or the variance avg over all $x$'s.

Prove for a single $x$ then average over all $x$'s.

Claim:

$$E_x[h(x)-p(x)]^2 = E_{x,y}[(h(x)-y)^2] - E_{x,y}[(p(x)-y)^2] \tag{20}$$

$$(h-p)^2 = E[(h-y)^2] - E[(p-y)^2] \tag{21}$$

$$(h-p)^2 = E[h^2 - 2hy + y^2] - E[p^2 - 2py + y^2] \tag{22}$$

$$(h-p)^2 = h^2 - 2h\underbrace{Ey}_{p} - p^2 + 2p\underbrace{Ey}_{p} = h^2 - 2hp + p^2 \tag{23}$$

$$(h-p)^2 = (h-p)^2 \tag{24}$$

Hence, we prove the claim from Equation 20 for a fixed $x$. To get the more general statement, we only need to average over random $x$. since

$$E_{x,y}[ANY] = E_x[E_y[ANY|x]] \tag{25}$$

# 3 Estimate $E[(h(x)-y)^2]$

We estimate $E[(h(x)-y)^2]$ by empirical average:

$$\hat{E}[(h(x)-y)^2] = \frac{1}{m}\sum (h(x_i)-y_i)^2 \tag{26}$$

$$L_h(x,y) = (h(x)-y)^2 \tag{27}$$

We want $E[L_h] \simeq \hat{E}[L_h]$ for all $h \in \mathcal{H}$

Chernoff bounds, union bound, VC-dim, growth function can all be generalized.

Q: How to minimize loss function for training set?

One answer: Perform a linear fit as shown in Figure 2.

Given $(x_1, y_1), \ldots, (x_m, y_m)$




Figure 2: Fit data from h(x) with a line.

$$min : \sum_i (wx_i - y_i)^2 \tag{28}$$

To minimize Equation 28 we set the derivative $\frac{d}{dw} = 2\sum_i (wx_i - y_i)x_i$ to 0 and get Equation 29:

$$w = \frac{\sum y_i x_u}{\sum x_i^2} \tag{29}$$

# 4 Generalize to more than one dimension

given $(\vec{x}_1, y_1), \ldots, (\vec{x}_m, y_m)$, $\vec{x}_i \in \mathcal{R}_n, y_i \in \mathcal{R}$

$\vec{w}$ using prediction rule $h(\vec{x}) = \vec{w} \cdot \vec{x}$

loss $(h) = \sum_i (\vec{w} \cdot \vec{x} - y_i)^2$

minimize: $\downarrow$

$$= \left|\left| \underbrace{\begin{pmatrix} \leftarrow \vec{x}_1^T \rightarrow \\ \leftarrow \vec{x}_2^T \rightarrow \\ \cdots \\ \leftarrow \vec{x}_m^T \rightarrow \end{pmatrix}}_{M} \underbrace{\begin{pmatrix} w_1 \\ w_2 \\ \cdots \\ w_m \end{pmatrix}}_{w} - \underbrace{\begin{pmatrix} y_1 \\ y_2 \\ \cdots \\ y_m \end{pmatrix}}_{b} \right|\right|_2^2$$

This can be solved by linear regression (next time).