

Parametric & Implicit Surfaces

Thomas Funkhouser
Princeton University
COS 426, Spring 2007

3D Object Representations

- Points
 - Range image
 - Point cloud
- Surfaces
 - Polygonal mesh
 - Subdivision
 - Parametric
 - Implicit
- Solids
 - Voxels
 - BSP tree
 - CSG
 - Sweep
- High-level structures
 - Scene graph
 - Application specific

Surfaces


- What makes a good surface representation?
 - Accurate
 - Concise
 - Intuitive specification
 - Local support
 - Affine invariant
 - Arbitrary topology
 - Guaranteed continuity
 - Natural parameterization
 - Efficient display
 - Efficient intersections



H&B Figure 10.46

Parametric Surfaces


- Boundary defined by parametric functions:
 - $x = f_x(u,v)$
 - $y = f_y(u,v)$
 - $z = f_z(u,v)$



FvDFH Figure 11.42

Parametric Surfaces


- Boundary defined by parametric functions:
 - $x = f_x(u,v)$
 - $y = f_y(u,v)$
 - $z = f_z(u,v)$



H&B Figure 10.10

Parametric Surfaces

- Example: surface of revolution
 - Take a curve and rotate it about an axis




Demetri Terzopoulos

Parametric Surfaces




- Example: swept surface
 - Sweep one curve along path of another curve



Parametric Surfaces



- How do we describe arbitrary smooth surfaces with parametric functions?




H&B Figure 10.46

Piecewise Polynomial Parametric Surfaces



- Surface is partitioned into parametric patches:




Same ideas as parametric splines!

Watt Figure 6.25

Parametric Patches



- Each patch is defined by blending control points




Same ideas as parametric curves!

FvDFH Figure 11.44

Parametric Patches



- Point $Q(u,v)$ on the patch is the tensor product of parametric curves defined by the control points




Watt Figure 6.21

Parametric Patches




- Point $Q(u,v)$ on the patch is the tensor product of parametric curves defined by the control points



Watt Figure 6.21

Parametric Patches


- Point $Q(u,v)$ on the patch is the tensor product of parametric curves defined by the control points



Watt Figure 6.21

Parametric Patches


- Point $Q(u,v)$ on the patch is the tensor product of parametric curves defined by the control points



Watt Figure 6.21

Parametric Patches

- Point $Q(u,v)$ on the patch is the tensor product of parametric curves defined by the control points



Watt Figure 6.21

Parametric Bicubic Patches

Point $Q(u,v)$ on any patch is defined by combining control points with polynomial blending functions:

$$Q(u,v) = \mathbf{U} \mathbf{M} \begin{bmatrix} P_{1,1} & P_{1,2} & P_{1,3} & P_{1,4} \\ P_{2,1} & P_{2,2} & P_{2,3} & P_{2,4} \\ P_{3,1} & P_{3,2} & P_{3,3} & P_{3,4} \\ P_{4,1} & P_{4,2} & P_{4,3} & P_{4,4} \end{bmatrix} \mathbf{M}^T \mathbf{V}^T$$


$$\mathbf{U} = [u^3 \ u^2 \ u \ 1] \quad \mathbf{V} = [v^3 \ v^2 \ v \ 1]$$

Where \mathbf{M} is a matrix describing the blending functions for a parametric cubic curve (e.g., Bezier, B-spline, etc.)

B-Spline Patches

$$Q(u,v) = \mathbf{U} \mathbf{M}_{\text{B-Spline}} \begin{bmatrix} P_{1,1} & P_{1,2} & P_{1,3} & P_{1,4} \\ P_{2,1} & P_{2,2} & P_{2,3} & P_{2,4} \\ P_{3,1} & P_{3,2} & P_{3,3} & P_{3,4} \\ P_{4,1} & P_{4,2} & P_{4,3} & P_{4,4} \end{bmatrix} \mathbf{M}_{\text{B-Spline}}^T \mathbf{V}$$

$$\mathbf{M}_{\text{B-Spline}} = \begin{bmatrix} -\frac{1}{6} & \frac{1}{2} & -\frac{1}{2} & \frac{1}{6} \\ \frac{1}{2} & -1 & \frac{1}{2} & 0 \\ -\frac{1}{2} & 0 & \frac{1}{2} & 0 \\ \frac{1}{6} & \frac{2}{3} & \frac{1}{6} & 0 \end{bmatrix}$$




Watt Figure 6.28

Bezier Patches

$$Q(u,v) = \mathbf{U} \mathbf{M}_{\text{Bezier}} \begin{bmatrix} P_{1,1} & P_{1,2} & P_{1,3} & P_{1,4} \\ P_{2,1} & P_{2,2} & P_{2,3} & P_{2,4} \\ P_{3,1} & P_{3,2} & P_{3,3} & P_{3,4} \\ P_{4,1} & P_{4,2} & P_{4,3} & P_{4,4} \end{bmatrix} \mathbf{M}_{\text{Bezier}}^T \mathbf{V}$$

$$\mathbf{M}_{\text{Bezier}} = \begin{bmatrix} -1 & 3 & -3 & 1 \\ 3 & -6 & 3 & 0 \\ -3 & 3 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix}$$




FDFH Figure 11.42

Bezier Patches



- Properties:
 - Interpolates four corner points
 - Convex hull
 - Local control




Watt Figure 6.22

Bezier Surfaces



- Continuity constraints are similar to the ones for Bezier splines




FvDFH Figure 11.43

Bezier Surfaces



- C^0 continuity requires aligning boundary curves




Watt Figure 6.26a

Bezier Surfaces



- C^1 continuity requires aligning boundary curves and derivatives




Watt Figure 6.26b

B-Spline Patches



$$Q(u, v) = \mathbf{U} \mathbf{M}_{\text{B-Spline}} \begin{bmatrix} P_{1,1} & P_{1,2} & P_{1,3} & P_{1,4} \\ P_{2,1} & P_{2,2} & P_{2,3} & P_{2,4} \\ P_{3,1} & P_{3,2} & P_{3,3} & P_{3,4} \\ P_{4,1} & P_{4,2} & P_{4,3} & P_{4,4} \end{bmatrix} \mathbf{M}_{\text{B-Spline}}^T \mathbf{V}$$

$$\mathbf{M}_{\text{B-Spline}} = \begin{bmatrix} -\frac{1}{6} & \frac{1}{2} & -\frac{1}{2} & \frac{1}{6} \\ \frac{1}{2} & -1 & \frac{1}{2} & 0 \\ -\frac{1}{2} & 0 & \frac{1}{2} & 0 \\ \frac{1}{6} & \frac{2}{3} & \frac{1}{6} & 0 \end{bmatrix}$$



Watt Figure 6.28

Parametric Surfaces



- Advantages:
 - Easy to enumerate points on surface
 - Possible to describe complex shapes
- Disadvantages:
 - Control mesh must be quadrilaterals
 - Continuity constraints difficult to maintain
 - Hard to find intersections


3D Object Representations

- Points
 - Range image
 - Point cloud
- Surfaces
 - Polygonal mesh
 - Subdivision
 - Parametric
 - Implicit
- Solids
 - Voxels
 - BSP tree
 - CSG
 - Sweep
- High-level structures
 - Scene graph
 - Application specific



Implicit Surfaces


- Represent surface with function over all space



Kazhdan

Implicit Surfaces

- Surface defined implicitly by function




Kazhdan



Implicit Surfaces


- Surface defined implicitly by function:
 - $f(x, y, z) = 0$ (on surface)
 - $f(x, y, z) < 0$ (inside)
 - $f(x, y, z) > 0$ (outside)



Turk

Implicit Surfaces

- Normals defined by partial derivatives
 - $\text{normal}(x, y, z) = (\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z})$



Normals



Tangents



Curvatures


Bloomenthal



Implicit Surface Properties

- (1) Efficient check for whether point is inside
 - Evaluate $f(x,y,z)$ to see if point is inside/outside/on

$$\left(\frac{x}{r_x}\right)^2 + \left(\frac{y}{r_y}\right)^2 + \left(\frac{z}{r_z}\right)^2 - 1 = 0$$



H&B Figure 10.10

Implicit Surface Properties

(2) Efficient surface intersections

- Substitute to find intersections

Ray: $P = P_0 + tV$

Sphere: $|P - O|^2 - r^2 = 0$

Substituting for P , we get:

$$|P_0 + tV - O|^2 - r^2 = 0$$

Solve quadratic equation:


$$at^2 + bt + c = 0$$

where:

$$a = 1$$

$$b = 2 V \cdot (P_0 - O)$$


$$c = |P_0 - O|^2 - r^2 = 0$$




Implicit Surface Properties

(3) Efficient boolean operations (CSG)

- Union, difference, intersect



Union




Difference

Bloomenthal

Implicit Surface Properties

(4) Efficient topology changes

- Surface is not represented explicitly!




Bourke

Implicit Surface Properties

(4) Efficient topology changes

- Surface is not represented explicitly!



Bloomenthal


Comparison to Parametric Surfaces

Implicit

- Efficient intersections & topology changes

Parametric

- Efficient “marching” along surface & rendering



Bloomenthal

Implicit Surface Representations

How do we define implicit function?

- Algebraic
- Blobby models
- Skeletons
- Procedural
- Samples
- Variational

Implicit Surface Representations



- How do we define implicit function?

- Algebraics
- Blobby models
- Skeletons
- Procedural
- Samples
- Variational


Algebraic Surfaces



- Implicit function is polynomial

$$f(x,y,z) = ax^d + by^d + cz^d + dx^{d-1}y + dx^{d-1}z + dy^{d-1}x + \dots$$

$$\left(\frac{x}{r_x}\right)^2 + \left(\frac{y}{r_y}\right)^2 + \left(\frac{z}{r_z}\right)^2 - 1 = 0$$



H&B Figure 10.10

Algebraic Surfaces



- Most common form: quadrics

$$f(x,y,z) = ax^2 + by^2 + cz^2 + 2dxy + 2eyz + 2fxz + 2gx + 2hy + 2jz + k$$

- Examples

- Sphere
- Ellipsoid
- Torus
- Paraboloid
- Hyperboloid




Menon


Algebraic Surfaces



- Higher degree algebraics



Cubic



Quartic




Degree six

Algebraic Surfaces



- Function extends to infinity
- Must trim to get desired patch (this is difficult!)




Algebraic Surfaces



- Equivalent parametric surface

- Tensor product patch of degree m and n curves yields algebraic function with degree $2mn$



Bicubic patch has degree 18!

Algebraic Surfaces



- Intersection
 - Intersection of degree m and n algebraic surfaces yields curve with degree mn



Intersection of bicubic patches has degree 324!

Implicit Surface Representations




- How do we define implicit function?
 - Algebraic
Blobby models
 - Skeletons
 - Procedural
 - Samples
 - Variational

Blobby Models




- Implicit function is sum of spherical basis functions



Blobby Models



- Sum of two blobs




Turk

Blobby Models



- Sum of four blobs



Turk

Blobby Models




- Blobby molecules

$$D(r) = ae^{-br^2}$$

- Meta balls

$$D(r) = \begin{cases} a(1 - \frac{3r^2}{b^2}) & 0 \leq r \leq b/3 \\ \frac{3a}{2}(1 - \frac{r}{b})^2 & b/3 \leq r \leq b \\ 0 & r \geq b \end{cases}$$




- Soft objects


$$D(r) = \begin{cases} a(1 - \frac{4r^6}{9b^6} + \frac{17r^4}{9b^4} - \frac{22r^2}{9b^2}) & r \leq b \\ 0 & r \geq b \end{cases}$$

Bourke

Blobby Model of Face




(a) $N = 1$




(b) $N = 2$

Blobby Model of Face




(c) $N = 10$




(d) $N = 35$

Blobby Model of Face




(e) $N = 70$




(f) $N = 243$

Blobby Model of Head




(a) $N = 1$




(b) $N = 2$

Blobby Model of Head




(c) $N = 20$




(d) $N = 60$

Blobby Model of Head



(e) $N = 120$



(f) $N = 451$

Blobby Models



Objects resulting from CSG of implicit soft objects and other primitives




Menon

Skeletons



- Bulge problem



Implicit Surface Representations




- How do we define implicit function?

- Algebraic
- Blobby models
- Skeletons
- Procedural
- Samples
- Variational

Skeletons



- Convolution surfaces



Bloomenthal

Implicit Surface Representations




- How do we define implicit function?

- Algebraic
- Blobby models
- Skeletons
- Procedural
- Samples
- Variational

Procedural Implicits



- $f(x,y,z)$ is result of procedure
 - Example: Mandelbrot set



H&B Figure 10.100

Implicit Surface Representations




- How do we define implicit function?
 - Algebraics
 - Blobby models
 - Skeletons
 - Procedural Samples
 - Variational

Sampled Functions



- Most common example: voxels
 - Interpolate samples stored on regular grid
 - Isosurface at $f(x,y,z) = 0$ defines surface



Sampled Functions



- Acquired from simulations or scans



Airflow Inside a Thunderstorm
(Bob Wilhelmson,
University of Illinois at Urbana-Champaign)




Visible Human
(National Library of Medicine)

Implicit Surface Representations




- How do we define implicit function?
 - Algebraics
 - Blobby models
 - Skeletons
 - Procedural Samples
 - Variational

Variational Implicit Surfaces



Turk

Example Implicit Surface



Turk

Implicit Surface Summary



- Advantages:
 - Easy to test if point is on surface
 - Easy to compute intersections/unions/differences
 - Easy to handle topological changes
- Disadvantages:
 - Indirect specification of surface
 - Hard to describe sharp features
 - Hard to enumerate points on surface
 - » Slow rendering

Summary



Feature	Polygonal Mesh	Implicit Surface	Parametric Surface	Subdivision Surface
Accurate	No	Yes	Yes	Yes
Concise	No	Yes	Yes	Yes
Intuitive specification	No	No	Yes	No
Local support	Yes	No	Yes	Yes
Affine invariant	Yes	Yes	Yes	Yes
Arbitrary topology	Yes	No	No	Yes
Guaranteed continuity	No	Yes	Yes	Yes
Natural parameterization	No	No	Yes	No
Efficient display	Yes	No	Yes	Yes
Efficient intersections	No	Yes	No	No