

# COS 511: Theoretical Machine Learning

Homework #2  
Sample size bounds and VC dimension

Due: February 28, 2008

---

## Problem 1

This problem explores another general method for bounding the error when the hypothesis space is infinite.

Some algorithms output hypotheses that can be represented by a small number of examples from the training set. For instance, suppose the domain is  $\mathbb{R}$  and we are learning a half-line of the form  $x \geq a$  where  $a$  defines the half-line. A simple algorithm chooses the left most positive training example  $a$  and outputs the corresponding half-line, which is clearly consistent with the data. Thus, in this case, the hypothesis can be represented by a single training example.

More formally, let  $F$  be a function mapping labeled examples to concepts, and assume that algorithm  $A$ , when given training examples  $(x_1, c(x_1)), \dots, (x_m, c(x_m))$  labeled by some unknown  $c \in \mathcal{C}$ , chooses some  $i_1, \dots, i_k \in \{1, \dots, m\}$  and outputs the consistent hypothesis  $F((x_{i_1}, c(x_{i_1})), \dots, (x_{i_k}, c(x_{i_k})))$ . In a sense, the algorithm has “compressed” the sample down to a sequence of just  $k$  of the  $m$  training examples.

- a. [5] Give such an algorithm for axis-aligned hyper-rectangles in  $\mathbb{R}^n$  with  $k = O(n)$ . (An axis-aligned hyper-rectangle is a set of the form  $[a_1, b_1] \times \dots \times [a_n, b_n]$ . For  $n = 2$ , this is the class of rectangles used repeatedly as an example in class.) Your algorithm should run in time polynomial in  $m$  and  $n$ .
- b. [15] Returning to the general case, assume as usual that the examples are chosen at random from some distribution  $D$ . Also assume that the size  $k$  is fixed. Argue *carefully* that the error of the output hypothesis  $h$ , with probability at least  $1 - \delta$  satisfies the bound:

$$\text{err}_D(h) \leq O\left(\frac{\ln(1/\delta) + k \ln m}{m - k}\right).$$

## Problem 2

[15] Let the domain be  $\mathbb{R}^d$ , and consider the class  $\mathcal{C}$  of linear threshold functions passing through the origin. That is, each such function is defined by a vector  $\mathbf{w} \in \mathbb{R}^d$  and is equal to 1 on points  $\mathbf{x}$  for which  $\mathbf{w} \cdot \mathbf{x} \geq 0$ , and 0 on all other points. Show that the VC-dimension of  $\mathcal{C}$  is exactly equal to  $d$ .

## Problem 3

[15] For each  $d = 0, 1, 2, \dots$ , give an example of a class  $\mathcal{C}$  for which Sauer’s Lemma is tight, i.e., for which the VC-dimension of  $\mathcal{C}$  is  $d$ , and, for each  $m$ ,  $\Pi_{\mathcal{C}}(m) = \sum_{i=0}^d \binom{m}{i}$ .