



Parametric Surfaces

COS 426, Spring 2022

Felix Heide

Princeton University



3D Object Representations

- Points
 - Range image
 - Point cloud
- Surfaces
 - Polygonal mesh
 - Parametric
 - Subdivision
 - Implicit
- Solids
 - Voxels
 - BSP tree
 - CSG
 - Sweep
- High-level structures
 - Scene graph
 - Application specific



Parametric Surfaces

- Applications
 - Design of smooth surfaces in cars, ships, etc.

Visualization





Parametric Surfaces


- Applications
 - Partitioning into surface patches.





Parametric Surfaces


- Applications
 - Design of smooth surfaces in cars, ships, etc.
 - Creating characters or scenes for movies





Parametric Curves


- Applications
 - Defining motion trajectories for objects or cameras





Parametric Curves

- Applications
 - Defining motion trajectories for objects or cameras
 - Defining smooth interpolations of sparse data



Google
Street View



Outline

- Parametric curves
 - Cubic B-Spline
 - Cubic Bézier
- Parametric surfaces
 - Bi-cubic B-Spline
 - Bi-cubic Bézier



Outline

- Parametric curves
 - Cubic B-Spline
 - Cubic Bézier
- Parametric surfaces
 - Bi-cubic B-Spline
 - Bi-cubic Bézier




Parametric Curves

- Defined by parametric functions:

- $x = f_x(u)$
- $y = f_y(u)$

- Example: line segment





Parametric Curves

- Defined by parametric functions:


- $x = f_x(u)$
- $y = f_y(u)$

- Example: line segment

$$f_x(u) = (1-u)x_0 + ux_1$$

$$f_y(u) = (1-u)y_0 + uy_1$$

$$u \in [0..1]$$



H&B Figure 10.10



Parametric Curves

- Defined by parametric functions:


- $x = f_x(u)$
- $y = f_y(u)$

- Example: circle

$$f_x(u) = r \cos(2\pi u)$$

$$f_y(u) = r \sin(2\pi u)$$

$$u \in [0..1]$$




H&B Figure 10.10



Parametric Curves

- Defined by parametric functions:

- $x = f_x(u)$
- $y = f_y(u)$



- Example: ellipse

$$f_x(u) = r_x \cos(2\rho u)$$

$$f_y(u) = r_y \sin(2\rho u)$$

$$u \hat{=} [0..1]$$

H&B Figure 10.10




Parametric curves

How to easily define arbitrary curves?

$$x = f_x(u)$$

$$y = f_y(u)$$






Parametric curves

How to easily define arbitrary curves?

$$x = f_x(u)$$

$$y = f_y(u)$$






Parametric curves


How to easily define arbitrary curves?

$$x = f_x(u)$$

$$y = f_y(u)$$



Slightly different notation than before:






Parametric curves

How to easily define arbitrary curves?

$$x = f_x(u)$$

$$y = f_y(u)$$



Use functions that “blend” control points

$$x = f_x(u) = V_{0x} * (1 - u) + V_{1x} * u$$

$$y = f_y(u) = V_{0y} * (1 - u) + V_{1y} * u$$

Simple functions of u




Parametric curves

More generally:

$$x(u) = \sum_{i=0}^n B_i(u) * Vi_x$$

$$y(u) = \sum_{i=0}^n B_i(u) * Vi_y$$






Parametric curves

More generally:

$$x(u) = \sum_{i=0}^n B_i(u) * Vi_x$$

$$y(u) = \sum_{i=0}^n B_i(u) * Vi_y$$

May use ***all*** points!





Parametric curves

What $B(u)$ functions should we use?

$$x(u) = \sum_{i=0}^n B_i(u) * Vi_x$$

$$y(u) = \sum_{i=0}^n B_i(u) * Vi_y$$




Parametric curves

What $B(u)$ functions should we use?

$$x(u) = \sum_{i=0}^n B_i(u) * Vi_x$$

$$y(u) = \sum_{i=0}^n B_i(u) * Vi_y$$








Parametric curves

What $B(u)$ functions should we use?

$$x(u) = \sum_{i=0}^n B_i(u) * Vi_x$$

$$y(u) = \sum_{i=0}^n B_i(u) * Vi_y$$






Parametric curves

What $B(u)$ functions should we use?

$$x(u) = \sum_{i=0}^n B_i(u) * Vi_x$$

$$y(u) = \sum_{i=0}^n B_i(u) * Vi_y$$









Parametric curves

What $B(u)$ functions should we use?

$$x(u) = \sum_{i=0}^n B_i(u) * Vi_x$$


$$y(u) = \sum_{i=0}^n B_i(u) * Vi_y$$





Blending Functions B

- Polynomial blending functions:



$$x(u) = \sum_{i=0}^n B_i(u) * Vi_x$$

$$y(u) = \sum_{i=0}^n B_i(u) * Vi_y$$




Blending Functions B

- Polynomial blending functions:

$$B_i(u) = \sum_{j=0}^m a_j u^j$$

$$x(u) = \sum_{i=0}^n B_i(u) * Vi_x$$

$$y(u) = \sum_{i=0}^n B_i(u) * Vi_y$$






Blending Functions B

- Polynomial blending functions:

$$B_i(u) = \sum_{j=0}^m a_j u^j$$

$$x(u) = \sum_{i=0}^n B_i(u) * V_i_x$$

$$y(u) = \sum_{i=0}^n B_i(u) * V_i_y$$




- Advantages of polynomials
 - Easy to compute
 - Easy to derive curve properties




Parametric Polynomial Curves

- Polynomial blending functions:

$$B_i(u) = \sum_{j=0}^m a_j u^j$$




- What degree polynomial?
 - Easy to compute
 - Easy to control
 - Expressive





Piecewise Parametric Polynomial Curves


- Splines:
 - Split curve into segments
 - Each segment defined by low-order polynomial blending subset of control vertices





Piecewise Parametric Polynomial Curves

- **Splines:**
 - Split curve into segments
 - Each segment defined by low-order polynomial blending subset of control vertices
- **Motivation:**
 - Same blending functions for every segment
 - Prove properties from blending functions
 - Provides **local control & efficiency**
- **Challenges**
 - How choose blending functions?
 - How determine properties?






Cubic Splines

- Some properties we might like to have:
 - Local control
 - Continuity
 - Interpolation?
 - Convex hull?

Blending functions determine properties

Properties determine blending functions

$$B_i(u) = \sum_{j=0}^m a_j u^j$$






Outline

- Parametric curves
 - Cubic B-Spline
 - Cubic Bézier
- Parametric surfaces
 - Bi-cubic B-Spline
 - Bi-cubic Bézier



Cubic B-Splines


- Properties:
 - Local control
 - C^2 continuity at joints
(infinitely continuous within each piece)
 - Approximating
 - Convex hull






Cubic B-Splines

Notes on continuity




[from Robert Duvall]

Geometric Continuity:



Parametric Continuity:




[from Carlo Séquin]




Cubic B-Splines


Notes on local control:




and convex hull property:




YES



YES



YES



NO

[from Steve Marschner]





Cubic B-Spline Blending Functions

Cubic B-Spline Curve Formation:

$$x(u) = \sum_{i=0}^n B_i(u) * V_i_x$$

$$y(u) = \sum_{i=0}^n B_i(u) * V_i_y$$

$$B_i(u) = \sum_{j=0}^m a_j u^j$$










Cubic B-Spline Blending Functions

In plot form backwards (start at 3):

$$B_i(u) = \sum_{j=0}^m a_j u^j$$





$$y_p(u) = \sum_{i=0}^n B_{-i}(u) * V_{p-i_x}$$

$$x_p(u) = \sum_{i=0}^n B_{-i}(u) * V_{p-i_y}$$



Cubic B-Spline Blending Functions



- A single blending function
- Local support
- Approximating





Cubic B-Spline Blending Functions

- Replicate blending functions
- Still local support





Try online at <http://bl.ocks.org/mbostock/4342190>



Cubic B-Spline Blending Functions



- How derive blending functions?
 - Cubic polynomials
 - Local control
 - C^2 continuity
 - Convex hull





Cubic B-Spline Blending Functions

- How derive blending functions?
 - Cubic polynomials
 - Local control
 - C^2 continuity
 - Convex hull






Cubic B-Spline Blending Functions

- Four cubic polynomials for four vertices
 - 16 variables (degrees of freedom)
 - Variables are a_i , b_i , c_i , d_i for four blending functions


$$\begin{aligned}b_{-0}(u) &= a_0 u^3 + b_0 u^2 + c_0 u^1 + d_0 \\b_{-1}(u) &= a_1 u^3 + b_1 u^2 + c_1 u^1 + d_1 \\b_{-2}(u) &= a_2 u^3 + b_2 u^2 + c_2 u^1 + d_2 \\b_{-3}(u) &= a_3 u^3 + b_3 u^2 + c_3 u^1 + d_3\end{aligned}$$





Cubic B-Spline Blending Functions

- C^2 continuity implies 15 constraints
 - Position of two curves same
 - Derivative of two curves same
 - Second derivatives same





Cubic B-Spline Blending Functions

Fifteen continuity constraints:

$$0 = b_{-0}(0)$$

$$0 = b_{-0}'(0)$$

$$0 = b_{-0}''(0)$$

$$b_{-0}(1) = b_{-1}(0)$$

$$b_{-0}'(1) = b_{-1}'(0)$$

$$b_{-0}''(1) = b_{-1}''(0)$$

$$b_{-1}(1) = b_{-2}(0)$$

$$b_{-1}'(1) = b_{-2}'(0)$$

$$b_{-1}''(1) = b_{-2}''(0)$$

$$b_{-2}(1) = b_{-3}(0)$$

$$b_{-2}'(1) = b_{-3}'(0)$$

$$b_{-2}''(1) = b_{-3}''(0)$$

$$b_{-3}(1) = 0$$

$$b_{-3}'(1) = 0$$

$$b_{-3}''(1) = 0$$

One more convenient constraint:

$$b_{-0}(0) + b_{-1}(0) + b_{-2}(0) + b_{-3}(0) = 1$$



Cubic B-Spline Blending Functions





- Solving the system of equations yields:

$$b_{-3}(u) = -\frac{1}{6}u^3 + \frac{1}{2}u^2 - \frac{1}{2}u + \frac{1}{6}$$

$$b_{-2}(u) = \frac{1}{2}u^3 - u^2 + \frac{2}{3}$$

$$b_{-1}(u) = -\frac{1}{2}u^3 + \frac{1}{2}u^2 + \frac{1}{2}u + \frac{1}{6}$$

$$b_{-0}(u) = \frac{1}{6}u^3$$





Cubic B-Spline Blending Functions

- In matrix form:

$$Q(u) = \sum_{i=0}^n B_i(u) * V_i \quad B_i(u) = \sum_{j=0}^m a_j u^j$$




Cubic B-Spline Blending Functions

- In matrix form:

$$Q(u) = \sum_{i=0}^n B_i(u) * V_i \quad B_i(u) = \sum_{j=0}^m a_j u^j$$

}

$$Q(u) = \begin{pmatrix} u^3 & u^2 & u & 1 \end{pmatrix} \frac{1}{6} \begin{pmatrix} -1 & 3 & -3 & 1 \\ 3 & -6 & 3 & 0 \\ -3 & 0 & 3 & 0 \\ 1 & 4 & 1 & 0 \end{pmatrix} \begin{pmatrix} V_0 \\ V_1 \\ V_2 \\ V_3 \end{pmatrix}$$



Outline


- Parametric curves
 - Cubic B-Spline
 - Cubic Bézier
- Parametric surfaces
 - Bi-cubic B-Spline
 - Bi-cubic Bézier



Bézier Curves

- Developed around 1960 by both
 - Pierre Bézier (Renault)
 - Paul de Casteljau (Citroen)
- Today: graphic design (e.g. FONTS)
- Properties:
 - Local control
 - Continuity **depends** on control points
 - **Interpolating** (every third for cubic)

Blending functions determine properties








Cubic Bézier Curves

Blending functions:

$$Q(u) = \sum_{i=0}^n B_i(u) * V_i \quad B_i(u) = \sum_{j=0}^m a_j u^j$$





Basic properties of Bézier Curves

- Endpoint interpolation:

$$Q(0) = V_0$$

$$Q(1) = V_n$$

- Convex hull:
 - Curve is contained within convex hull of control polygon
- Symmetry

$$Q(u) \text{ defined by } \{V_0, \dots, V_n\} \equiv Q(1-u) \text{ defined by } \{V_n, \dots, V_0\}$$



Cubic Bézier Curves

Bézier curves in matrix form:

$$\begin{aligned} Q(u) &= \sum_{i=0}^n V_i \binom{n}{i} u^i (1-u)^{n-i} \\ &= (1-u)^3 V_0 + 3u(1-u)^2 V_1 + 3u^2(1-u) V_2 + u^3 V_3 \end{aligned}$$


$$= \begin{pmatrix} u^3 & u^2 & u & 1 \end{pmatrix} \begin{pmatrix} -1 & 3 & -3 & 1 \\ 3 & -6 & 3 & 0 \\ -3 & 3 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} V_0 \\ V_1 \\ V_2 \\ V_3 \end{pmatrix}$$

$\mathbf{M}_{\text{Bézier}}$



Bézier Curves

- Curve $Q(u)$ can also be defined by nested interpolation:




V_i are *control points*

$\{V_0, V_1, \dots, V_n\}$ is *control polygon*



Bézier Curves

- Curve $Q(u)$ can also be defined by nested interpolation:




V_i are *control points*

$\{V_0, V_1, \dots, V_n\}$ is *control polygon*



Bézier Curves

- Curve $Q(u)$ can also be defined by nested interpolation:




V_i are *control points*

$\{V_0, V_1, \dots, V_n\}$ is *control polygon*



Bézier Curves

- Curve $Q(u)$ can also be defined by nested interpolation:




V_i are *control points*

$\{V_0, V_1, \dots, V_n\}$ is *control polygon*



Bézier Curves

- Curve $Q(u)$ can also be defined by nested interpolation:




V_i are *control points*

$\{V_0, V_1, \dots, V_n\}$ is *control polygon*



Bézier Curves

- Curve $Q(u)$ can also be defined by nested interpolation:




V_i are *control points*

$\{V_0, V_1, \dots, V_n\}$ is *control polygon*



Enforcing Bézier Curve Continuity

- C⁰: $V_3 = V_4$
- C¹: $V_5 - V_4 = V_3 - V_2$
- C²: $V_6 - 2V_5 + V_4 = V_3 - 2V_2 + V_1$





Outline


- Parametric curves
 - Cubic B-Spline
 - Cubic Bézier
- Parametric surfaces
 - Bi-cubic B-Spline
 - Bi-cubic Bézier




Parametric Surfaces

- Defined by parametric functions:

- $x = f_x(u, v)$
- $y = f_y(u, v)$
- $z = f_z(u, v)$



Parametric functions
define mapping from
(u, v) to (x, y, z):



FvDFH Figure 11.42




Parametric Surfaces

- Defined by parametric functions:

- $x = f_x(u, v)$
- $y = f_y(u, v)$
- $z = f_z(u, v)$

- Example: quadrilateral



$$f_x(u, v) = (1 - v)((1 - u)x_0 + ux_1) + v((1 - u)x_2 + ux_3)$$

$$f_y(u, v) = (1 - v)((1 - u)y_0 + uy_1) + v((1 - u)y_2 + uy_3)$$

$$f_z(u, v) = (1 - v)((1 - u)z_0 + uz_1) + v((1 - u)z_2 + uz_3)$$




Parametric Surfaces

- Defined by parametric functions:

- $x = f_x(u, v)$
- $y = f_y(u, v)$
- $z = f_z(u, v)$

- Example: quadrilateral



$$f_x(u, v) = (1 - v)((1 - u)x_0 + ux_1) + v((1 - u)x_2 + ux_3)$$

$$f_y(u, v) = (1 - v)((1 - u)y_0 + uy_1) + v((1 - u)y_2 + uy_3)$$

$$f_z(u, v) = (1 - v)((1 - u)z_0 + uz_1) + v((1 - u)z_2 + uz_3)$$



Parametric Surfaces

- Defined by parametric functions:


- $x = f_x(u, v)$
- $y = f_y(u, v)$
- $z = f_z(u, v)$

- Example: ellipsoid

$$f_x(u, v) = r_x \cos v \cos u$$

$$f_y(u, v) = r_y \cos v \sin u$$

$$f_z(u, v) = r_z \sin v$$




H&B Figure 10.10



Parametric Surfaces


To model arbitrary shapes, surface is partitioned into parametric patches





Parametric Patches

- Each patch is defined by blending control points




FvDFH Figure 11.44



Parametric Patches

- Each patch is defined by blending control points





Same ideas as parametric curves!

FvDFH Figure 11.44



Parametric Patches

- Point $Q(u,v)$ on the patch is the tensor product of parametric curves defined by the control points





Watt Figure 6.21



Parametric Patches

- Point $Q(u,v)$ on the patch is the tensor product of parametric curves defined by the control points





Watt Figure 6.21



Parametric Patches

- Point $Q(u,v)$ on the patch is the tensor product of parametric curves defined by the control points





Watt Figure 6.21



Parametric Patches

- Point $Q(u,v)$ on the patch is the tensor product of parametric curves defined by the control points





Watt Figure 6.21



Parametric Patches

- Point $Q(u,v)$ on the patch is the tensor product of parametric curves defined by the control points



Watt Figure 6.21



Parametric Bicubic Patches

Point $Q(u,v)$ on any patch is defined by combining control points with polynomial blending functions:

$$Q(u, v) = \mathbf{U} \mathbf{M} \begin{bmatrix} P_{1,1} & P_{1,2} & P_{1,3} & P_{1,4} \\ P_{2,1} & P_{2,2} & P_{2,3} & P_{2,4} \\ P_{3,1} & P_{3,2} & P_{3,3} & P_{3,4} \\ P_{4,1} & P_{4,2} & P_{4,3} & P_{4,4} \end{bmatrix} \mathbf{M}^T \mathbf{V}^T$$

$$\mathbf{U} = [u^3 \quad u^2 \quad u \quad 1] \quad \mathbf{V} = [v^3 \quad v^2 \quad v \quad 1]$$


Where \mathbf{M} is a matrix describing the blending functions for a parametric cubic curve (e.g., Bézier, B-spline, etc.)



B-Spline Patches

$$Q(u, v) = \mathbf{U} \mathbf{M}_{\text{B-Spline}} \begin{bmatrix} P_{1,1} & P_{1,2} & P_{1,3} & P_{1,4} \\ P_{2,1} & P_{2,2} & P_{2,3} & P_{2,4} \\ P_{3,1} & P_{3,2} & P_{3,3} & P_{3,4} \\ P_{4,1} & P_{4,2} & P_{4,3} & P_{4,4} \end{bmatrix} \mathbf{M}_{\text{B-Spline}}^T \mathbf{V}$$

$$\mathbf{M}_{\text{B-Spline}} = \begin{bmatrix} -1/6 & 1/2 & -1/2 & 1/6 \\ 1/2 & -1 & 1/2 & 0 \\ -1/2 & 0 & 1/2 & 0 \\ 1/6 & 2/3 & 1/6 & 0 \end{bmatrix}$$




Watt Figure 6.28



Bézier Patches

- Properties:
 - Interpolates four corner points
 - Convex hull
 - Local control




Watt Figure 6.22



Bézier Patches

$$Q(u, v) = \mathbf{U} \mathbf{M}_{\text{Bezier}} \begin{bmatrix} P_{1,1} & P_{1,2} & P_{1,3} & P_{1,4} \\ P_{2,1} & P_{2,2} & P_{2,3} & P_{2,4} \\ P_{3,1} & P_{3,2} & P_{3,3} & P_{3,4} \\ P_{4,1} & P_{4,2} & P_{4,3} & P_{4,4} \end{bmatrix} \mathbf{M}_{\text{Bezier}}^T \mathbf{V}$$

$$\mathbf{M}_{\text{Bezier}} = \begin{bmatrix} -1 & 3 & -3 & 1 \\ 3 & -6 & 3 & 0 \\ -3 & 3 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix}$$




FvDFH Figure 11.42



Piecewise Polynomial Parametric Surfaces


Surface is composition of many parametric patches





Piecewise Polynomial Parametric Surfaces

Must maintain continuity across seams




Same ideas as parametric splines!

Watt Figure 6.25



Bézier Surfaces

- Continuity constraints are similar to the ones for Bézier splines




FvDFH Figure 11.43




Bézier Surfaces

- C^0 continuity requires aligning boundary curves



Control point polyhedra



Patches


Watt Figure 6.26a




Bézier Surfaces

- C^1 continuity requires aligning boundary curves and derivatives

Four sets of three control points must be collinear



Control point polyhedra



Patches

Watt Figure 6.26b



Parametric Surfaces

- Properties
 - ? Natural parameterization
 - ? Guaranteed smoothness
 - ? Intuitive editing
 - ? Concise
 - ? Accurate
 - ? Efficient display
 - ? Easy acquisition
 - ? Efficient intersections
 - ? Guaranteed validity
 - ? Arbitrary topology



AUTODESK.



Parametric Surfaces

- Properties
 - ☺ Natural parameterization
 - ☺ Guaranteed smoothness
 - ☺ Intuitive editing
 - ☺ Concise
 - ☺ Accurate
 - Efficient display
 - ☹ Easy acquisition
 - ☹ Efficient intersections
 - ☹ Guaranteed validity
 - ☹ Arbitrary topology

